

ANIMAL HEALTH
PRODUCTS



ipm



Index

ANTIBACTERIAL

Injectable Products

Flormeg Solution for Injection.....	6
Florsym Solution for Injection.....	7
Linkosym Solution for Injection.....	8
Marboforce Solution for Injection.....	9
Oksipirimavet LA Solution for Injection.....	10
Pentril %10 Solution for Injection.....	11
Ribogen Solution for Injection.....	12
Sulfakill Solution for Injection.....	13
Symtil 300 Solution for Injection.....	14
Taylobar Solution for Injection.....	15
Tulasym Solution for Injection.....	16

Oral Products

Doksisym %50 Water Soluble Powder.....	18
Doksol %10 Oral Solution.....	19
Enrosym %10 Oral Solution.....	20
Enrosym %20 Oral Solution.....	21
Enrosym %80 Water Soluble Powder.....	22
Erizol Water Soluble Powder.....	23
Florsym %30 Oral Solution.....	24
Kolisym %24 Water Soluble Powder.....	25
Linkosym Water Soluble Powder.....	26
Metrisym Oral Tablet.....	27
Neosym %50 Water Soluble Powder.....	28
Paripto Powder for Oral Solution.....	29
Sulfasym Oral Suspension.....	30
Symtay Water Soluble Powder.....	31
Symtil %30 Oral Solution.....	32
Taylosym %24 Oral Solution.....	33
Tetrasym %80 Water Soluble Powder.....	34

ANTIPARASITIC

Injectable Products

Kantel Solution for Injection.....	35
Midokar Solution for Injection.....	36
Klosym Solution for Injection.....	37
Symec %1 Solution for Injection.....	38
Symec Fascio Solution for Injection.....	39

Oral Products

Albesym %10 Oral Suspension.....	40
Albesym 1200 Oral Tablet.....	41
Medizanid Oral Tablet.....	42
Wormkill Oral Tablet.....	43

Topical Solutions

Flumetsym Pour on Solution.....	44
---------------------------------	----

ANTICOCCIDIAL

Amprolsym %25 Oral Solution.....	45
Symcox Oral Solution.....	46

CLINICAL MEDICATIONS

IPM Ksilazin Solution for Injection.....	47
Furosym Solution for Injection.....	48
Symatro Solution for Injection.....	49
Symbuton Solution for Injection.....	53

VITAMIN

Adecure AD3E Solution for Injection.....	50
Adezek Solution for Injection.....	51
Bevitbar Solution for Injection.....	52
Symvit-C Solution for Injection.....	54
Symfos Solution for Injection.....	55

MINERAL

Symcal Solution for Injection.....	56
------------------------------------	----

ANALGESIC- ANTI-INFLAMMATORY

Badeljin Solution for Injection.....	57
Ketozek Solution for Injection.....	58
Melosym Solution for Injection.....	59
Melosym X8 Solution for Injection.....	60
Symful Solution for Injection.....	61



ANIMAL HEALTH PRODUCTS

ipm

Flormeg

Solution for Injection

Flormeg Solution for Injection
300 mg florfenicol, 16.5 mg flunixin

TARGET SPECIES: cattle

INDICATIONS: It is used in cattle for the treatment of respiratory infections caused by *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis* and *Histophilus somni*.

PACKAGING: 100 ml, 250 ml

DOSE: 2 ml/15 kg bw/day, one application subcutaneously

WITHDRAWAL TIME: Meat: 46 days.

Milk: Not authorized for use in lactating animals producing milk for human consumption.

See the leaflet for detailed information.



Florsym

Solution for Injection

Florsym Solution for Injection
300 mg florfenicol

TARGET SPECIES: cattle, sheep

INDICATIONS:**Cattle:** Treatment and metaphylaxis of respiratory tract infections caused by florfenicol-susceptible strains of Mannheimia haemolytica, Pasteurella multocida and Histophilus somni. Before metaphylactic use, the presence of disease in the flock should be ensured. **Sheep:** respiratory tract infections from Mannheimia haemolytica and Pasteurella multocida.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml/ 15 kg b.w. (i.m.),
2 ml/ 15 kg b.w.(s.c. for cattle only) treatment should continue at 48 hours intervals.

Sheep: 1 ml/15 kg bw (i.m.) for 3 days

WITHDRAWAL TIME:Meat: 30 days (i.m., cattle), 39 days (sheep, 44 days (s.c., cattle) Do not use in lactating cattle producing milk (including dry period) for human consumption

See the leaflet for detailed information.



Linkosym

Solution for Injection

Linkosym Solution for Injection

Lincomycin hydrochloride (equivalent to 222 mg lincomycin base)
Spectinomycin sulfate tetrahydrate (equivalent to 444 mg spectinomycin base)

TARGET SPECIES:sheep, lambs, calves, cattle, dog, cat, broiler

INDICATIONS:It is used in the treatment of respiratory, digestive, urinary system infections and soft tissue infections and arthritis caused by bacteria sensitive to lincomycin and spectinomycin in target species.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Sheep-goat: 1 ml for 10 kg bodyweight during 3-5 days

Calf -Steer, Cattle: 1 ml for 10 kg bodyweight, intramuscularly,daily during 3-5 days(first day 2 injections)

Dog-Cat: 1 ml for 5 kg body weight 1-2 a day, during 3-7 days.

Chikens and Turkey: 1 ml for 5 kg, subcutaneously for 3 days

WITHDRAWAL TIME:

Meat: 30 days for Cattle,Sheep ve Chicken

Milk: 72 hours for cows

Not to be used in sheep producing milk intended for human consumption

Not to be used in chicken producing egg intended for human consumption

See the leaflet for detailed information.



Marboforce

Solution for Injection

Marboforce Solution for Injection
100 mg marbofloxacin

TARGET SPECIES: cattle

INDICATIONS:It is indicated for respiratory tract infections caused by *Pasteurella multocida*, *Mannheimia haemolytica* ve *Histophilus somni* and used for the treatment of acute mastitis and infections caused by *Escherichia coli* and *Mycoplasma bovis*.

PACKAGING:100 ml

DOSE: For respiratory track infections dose is 2 ml/25 kg b.w. (i.m.). For *Mycoplasma bovis* infections and acute mastitis dose is 1 ml/50kg b.w./ day (i.m.,s.c.), should be applied for 3 days.

WITHDRAWAL TIME:

Meat: 3 days (for 8 mg/ kg b.w. treatment), 6 days (for 2 mg/ kg b.w. treatment)

Milk: 3 days (6 milkings, 8 mg/ kg b.w.), 2 days (4 milkings, 2 mg/ kg b.w.)

See the leaflet for detailed information.



Oksipirimavet LA

Solution for Injection

Oksipirimavet LA Solution for Injection
200 mg oxytetracycline

TARGET SPECIES: cattle, sheep, goat

INDICATIONS: It indicated to treatment of systemic, regional and respiratory system infections caused by bacteria sensitive to oxytetracycline.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 10 kg/bw (i.m)
If needed the dose is repeated after 4-5 days

WITHDRAWAL TIME:

Meat: 28 days

Milk: 10 days (cattle), 7 days (sheep, goat)

See the leaflet for detailed information.



Pentril %10

Solution for Injection

Pentril %10 Solution for Injection
100 mg Enrofloxacin

TARGET SPECIES:cattle, sheep, goat

INDICATIONS:It is used for the treatment of sensitive gram negative, gram positive and Mycoplasma related infections in cattle and sheep and goats.

PACKAGING: 20 ml, 50 ml, 100 ml, 250 ml

DOSE: Practical dose: 1 ml for 20 kg bw/day Product may given subcutaneously or intravenously. Treatment should continue for 3-5 days.

WITHDRAWAL TIME:

Meat: 5 days (cattle i.v.), 12 days (cattle s.c.), 4 days (sheep), 6 days (goats)

Milk: 3 days (sheep, cattle i.v.), 4 days (goats, cattle s.c.)

See the leaflet for detailed information.



Ribogen

Solution for Injection

Ribogen Solution for Injection
100 mg Gentamycine

TARGET SPECIES: cattle, horse, dog, cat

INDICATIONS: It is used against gentamicin sensitive bacteria causing respiratory, digestive and urogenital system infections, septicaemia and other soft tissue infections treatment in cattle, horse, dog and cat.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Cattle: 2 ml / 100kg body weight for 3-7 days

Horse: 6.5 ml/100kg bw for 3-5 days

Dog and cat: 0.2ml / 5 kg body weight for 3-5 days

WITHDRAWAL TIME:Meat: 139 days(cattle), 103 days(calves) for 3 injections,

192 days for 6 injections, 214 days for 7 injections

Milk: 7 days

See the leaflet for detailed information.



Sulfakill

Solution for Injection

Sulfakill Oral Suspension
200 mg sulfadimidin, 40 mg trimetoprim

TARGET SPECIES: Cattle, Horse, Cat, Dog

INDICATIONS: Sulfakill Oral Solution is indicated for the infections caused by susceptible microorganisms

PACKAGING: 50 ml, 100 ml ve 250 ml

DOSE: 1 ml/10-15 kg b.w. for cattle and horses; 1 ml/15 kg bw for cats and dogs.

WITHDRAWAL TIME: Meat: 12 days **Milk:** 5 days

See the leaflet for detailed information.



Symtil 300

Solution for Injection

Symtil 300 Solution for Injection
300 mg tilmicosin

TARGET SPECIES: cattle, sheep

INDICATIONS: SYMTIL 300 Solution for injection is indicated for treatment of respiratory disease (BRD) caused by bacteria susceptible to tilmicosin, treatment of respiratory disease associated with *Pasteurella haemolytica*, treatment of interdigital necrobacillosis and foot rot caused by *Fusobacterium necrophorus*, prevention of abortus caused by *Chlamidia psittaci* and as parenteral support in the treatment of mastitis and metritis in cattle and sheep.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 30 kg b.w.

WITHDRAWAL TIME:Meat: 70 days (cattle), 42 days (sheep) **Milk:** 18 days (sheep) 36 days(cattle)

See the leaflet for detailed information.



Taylobar

Solution for Injection

Taylobar Solution for Injection
200 mg tylosin

TARGET SPECIES: cattle, sheep, goat, cat, dog

INDICATIONS:Cattle (adult): metritis caused by gram positive bacteria, respiratory system infections, Streptococcus spp., Staphylococcus spp. Treatment of mastitis caused by Fusobacterium necrophorum and interdigital necrobacillosis caused by Fusobacterium necrophorum, Calf: Necrobacillosis (Fusobacterium necrophorum) and respiratory system infections **Sheep-goat:** metritis caused by gram positive bacteria, respiratory system infections, Gram positive bacteria or Mycoplasma spp. mastitis caused by Dog: Respiratory tract infections and otitis caused by susceptible bacteria Cat: respiratory tract infections caused by susceptible bacteria

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE:

Applied intramuscularly and only applied slow intravenous for cattle.

Cattle: 2.5-5 ml/100 kg bw

Sheep-goat: 2.5 ml/50 kg bw

Cat-dog: 0.35-0.55 ml/10 kg bw

WITHDRAWAL TIME:Meat: 28 days (cattle) 42 days (sheep, goat)

Milk: 108 hours

See the leaflet for detailed information.



Tulasym

Solution for Injection

Tulasym Solution for Injection
100 mg tulatromycin

TARGET SPECIES: cattle, sheep

INDICATIONS: Cattle; It is indicated for the treatment and metaphylaxis of bovine respiratory disease (BRD) caused by Mannheimia haemolytica, Pasteurella multocida, Histophilus somni and Mycoplasma bovis. It is indicated for the treatment of infectious keratoconjunctivitis caused by Morexella bovis sensitive to tulatromycin

Sheep; It is indicated for the early treatment of pododermatitis caused by virulent Dichelobacter nodosus

PACKAGING: 100 ml, 250 ml

DOSE: Cattle: 2,5 mg tulatromycin/kg b.w. (1ml/40 kg b.w.) single dose via s.c.

Sheep: 2,5 mg tulatromycin/kg b.w. (1ml/40 kg b.w.) single dose via i.m. route

WITHDRAWAL TIME:

Meat: Cattle: 22 days, **Sheep:** 16 days

Not to be used in animals producing milk intended for human consumption or animals expected to give birth in two months that will produce milk for human consumption.

See the leaflet for detailed information.





Doksisym %50

Water Soluble Powder

Doksisym %50 Water Soluble Powder
577 mg Doxycycline hyclate (equivalent to 500 mg base)

TARGET SPECIES: Non-ruminating Calves, Chicken, Turkey

INDICATIONS: It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

PACKAGING: 1 kg

DOSE:

Calves; 17.3 mg product/b.w/day via oral route during 3 or 5 days

Chicken; 43.3 mg product/b.w/day via oral route during 3 or 5 days

Turkey; 50.2 mg product/b.w/day via oral route during 3 or 5 days

WITHDRAWAL TIME:

Meat: 7 days (calves), 5 days (chicken), 12 days (turkey)

Egg: Do not use in animals producing egg for human consumption.

Milk: Do not use in animals producing milk for human consumption.

Not to be used within 4 weeks before egg production period.

See the leaflet for detailed information.



Doksol %10

Oral Solution

Doksol %10 Oral Solution
100 mg Doxycycline

TARGET SPECIES: Calves, Chicken, Turkey

INDICATIONS: It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

PACKAGING: 100 ml, 1 L

DOSE:

Recommended dose of product is mixed in water

Calves; 0.08 ml product/b.w/day via oral route during 3-5 days

Chicken; 0.21 ml product/b.w/day via oral route during 3-5 days

Turkey; 0.25 ml product/b.w/day via oral route during 3-5 days

WITHDRAWAL TIME:

Meat: 7 days (calves), 5 days (chicken), 12 days (turkey)

Egg: Do not use in animals producing egg for human consumption.

Milk: Do not use in animals producing milk for human consumption.

Not to be used within 4 weeks before egg production period.

See the leaflet for detailed information.



Enrosym %10

Oral Solution

Enrosym %10 Oral Solution
100 mg enrofloxacin

TARGET SPECIES: chicken, turkey, rabbit


INDICATIONS: It is indicated for the treatment of the infections caused by gram negative and positive bacteria and Mycoplasma spp. susceptible to enrofloxacin in broiler turkeys, chickens and rabbits.

PACKAGING: 100 ml, 2500 ml

DOSE: The daily dose to be administered is 10 mg/kg body weight by adding drinking water for 3-5 days (5 days for rabbit).

WITHDRAWAL TIME:Meat: 7 days (chicken), 13 days (turkey), 3 days (rabbit) Should not be given to chickens and turkeys producing eggs for human consumption, not to be used within 14 days to egg laying period.

See the leaflet for detailed information.



Enrosym %20

Oral Solution

Enrosym %20 Oral Solution
200 mg enrofloxacin

TARGET SPECIES: Chicken, Turkey, Rabbit

INDICATIONS: It is used in the treatment of respiratory and gastro intestinal system infections caused by susceptible bacteria in chicken, turkey and rabbits.


PACKAGING: 1 L, 3 L

DOSE: Practically 5 ml of Enrosym %20 Oral Solution per 100 kg body weight is added to drinking water daily. It is used for 3-5 days (5 days in rabbit)

WITHDRAWAL TIME: Meat: 7 days (chicken), 13 days (turkey), 3 days (rabbit)

Egg: Do not use in animals producing egg for human consumption. Not to be used within 14 days to egg laying period.

See the leaflet for detailed information.



Enrosym %80

Water Soluble Powder

Enrosym %80 Water Soluble Powder
800 mg enrofloxacin

TARGET SPECIES: chicken, turkey,

INDICATIONS: Enrosym 80% Oral Solution Powder is indicated for the treatment of infections caused by gram-negative and gram-positive bacteria that are enrofloxacin-sensitive, infections caused by mycoplasmas and secondary bacterial infections accompanying viral infections in broiler chickens and broiler turkeys. It is indicated mainly for the following: CRD (Chronic Respiratory Disease), mycoplasmosis, coliseptisemia, colibacillosis, pasteurellosis

PACKAGING: 1 kg

Dose: 1,25 g/100 kg total b.w./day

WITHDRAWAL TIME:Meat: 12 days (chicken), 14 days (turkey).

Not to be used in chicken and turkey producing egg intended for human consumption

See the leaflet for detailed information.



Erizol

Water Soluble Powder

Erizol Water Soluble Powder
509 mg Eritromycin

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Treatment of infections associated with Mycoplasma

PACKAGING: 100 g and 1 kg

DOSE: It is used as 20-40 mg/g erythromycine base/day via oral route during 3 or 5 days
Practically 39-79 g/1000 kg b.w.

WITHDRAWAL TIME:

Meat: 3 days (chicken), 6 days (turkey)

Egg: 5 days

See the leaflet for detailed information.



Florsym %30

Oral Solution

Florsym %30 Oral Solution
300 mg florfenicol

TARGET SPECIES: Chicken

INDICATIONS: Florsym 30% Oral Solution is indicated for the treatment of respiratory diseases, sacculitis and colisepticemia caused by E. Coli.

PACKAGING: 1000 ml and 2500 ml

DOSE: 100ml/1500 kg total b.w./ day

WITHDRAWAL TIME:

Meat: 5 days

Not to be used in chicken producing egg intended for human consumption

See the leaflet for detailed information.



Kolisym %24

Water Soluble Powder

Kolisym %24 Water Soluble Powder
4.800.000 IU kolistin

TARGET SPECIES: Chicken, Turkey, Calves, Lambs

INDICATIONS: It is indicated for the treatment of digestive system infections caused by Gram negative bacteria including especially E. Coli and Salmonella species in chicken, turkey, calves and lambs.

PACKAGING: 500-1000-2500 g

DOSE:

25g/1000 kg total b.w. for chicken and turkey 3-4 days

1 g/50 kg b.w. is sufficient in calves and lambs

WITHDRAWAL TIME:

Meat: 1 day

Egg: zero "0" days

not to be used on animals that produce milk for human consumption.

See the leaflet for detailed information.



Linkosym

Water Soluble Powder

Linkosym Water Soluble Powder
222 mg lincomycin, 444 mg spectinomycin

TARGET SPECIES: Broiler, Turkey

INDICATIONS: Linkosym Oral Solution Powder is indicated for the treatment of the infections caused by sensitive bacteria, chronic respiratory infections (CRD) caused especially by *Mycoplasma* sp., and CRD complex caused both by *Mycoplasma* sp. and coliform microorganisms, avian cholera and other coliform infections in broiler chickens and poults, as well as air sac infections in turkeys.

PACKAGING: 150 g

DOSE: 7.5kg/100 kg total b.w. for chicken and 22.5 kg/100 kg b.w. for turkey

WITHDRAWAL TIME:

Meat: Chickens 2, Turkey 8 days.

Not to be used in poultry producing egg intended for human consumption

See the leaflet for detailed information.



Metrisym

Oral Tablet

Metrisym Oral Tablet
1000 mg Sulfadiazin 200 mg Trimethoprim

TARGET SPECIES: calves

INDICATIONS: Metrisym Oral Tablet is mainly indicated for the treatment of bacterial diarrhea in calves, it can also used in the treatment of acute salmonellosis and bacterial pneumonia.

PACKAGING: 2x 5 tablets

DOSE:

1 tablet/ 40 kg b.w./ day
during 5 days

WITHDRAWAL TIME: Meat: 15 days

See the leaflet for detailed information.



Neosym %50

Water Soluble Powder

Neosym %50 Water Soluble Powder
500 mg neomycin

TARGET SPECIES: Calves, Lambs, Kids, Chicken, Turkey

INDICATIONS: It is indicated for the treatment of infections caused by neomycin sensitive bacteria in calves, lambs, kids chicken and turkey.

PACKAGING: 20 g, 100 g, 1 kg

DOSE: 1 gr/ 20 kg b.w./ day, split in 2 for lambs, calves and kids for 3-5 days

WITHDRAWAL TIME:

Meat: 14 days

Egg: zero "0" days

See the leaflet for detailed information.



Paripto

Powder for Oral Solution

100 mg paromomycine sulfate

TARGET SPECIES: non-ruminating calves

INDICATIONS: Paripto Powder for Oral Solution is used for the treatment of gastro-intestinal infections caused by E. Coli, Salmonella and Cryptosporidium in non-ruminating calves

PACKAGING: 1 kg, 2,5 kg

DOSE: Used by adding to drinking water
2,5-5 g/ 10 kg bw/day for Colibacillosis and Salmonellosis, during 3-5 days. 5 g/ 10 kg bw/day for Cryptosporidiosis during 7 days

WITHDRAWAL TIME: 62 days (5 g/ 10 kg b.w. for 7 days treatment)
Meat: 30 days (2,5-5 g/ 10 kg b.w. for 3-5 days treatment)

See the leaflet for detailed information.



Sulfasym

Oral Suspension

Sulfasym Oral Suspension
400 mg Sulphamethoxazole, 80 mg Trimetoprim

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Sulfasym is indicated for the treatment of respiratory infections caused by trimetoprim- and sulfamethoxazole-sensitive *Escherichia Coli* in avians (broiler chickens - turkeys).

PACKAGING: 100 ml, 1 L, 2.5 L

DOSE: There should be 1 ml Sulfasym Oral Suspension per 3.64 kg body weight. Treatment should be continued for 3-4 days

WITHDRAWAL TIME:Meat: 10 days Should not given to poultry producing eggs for human consumption

See the leaflet for detailed information.



Symtay

Symtay Water Soluble Powder
50 mg tylosin tartrate

TARGET SPECIES: chicken, turkey

INDICATIONS: SYMTAY Oral Solution Powder is indicated for the treatment of infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD), Leptospirosis and infectious sinus inflammation in turkeys.

Water Soluble Powder

PACKAGING: 20 g , 100-500-1000 and 2500g

DOSE: 5.5 g SYMTAY per 100 kg

WITHDRAWAL TIME:Meat: 2 days (chicken), 5 days (Turkey)

The chicken eggs obtained during the treatment and for a period of 5 days following the last drug administration should not be used for human consumption.

See the leaflet for detailed information.



Symtil %30

Oral Solution

Symtil %30 Oral Solution
300 mg tilmicosine

TARGET SPECIES: Broiler, Turkey, Calves

INDICATIONS: SYMTAY Oral Solution Powder is indicated for the treatment of infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD), Leptospirosis and infectious sinus inflammation in turkeys.

PACKAGING: 250-500, 1000 ml.

DOSE: 5-7 ml/ 100 kg total b.w.

WITHDRAWAL TIME: Meat: 12 days (chicken), 19 days (turkey), 42 days (calf)

Not to be used in chicken and turkey producing egg intended for human consumption. Not to be used within 14 days of egg laying period.

See the leaflet for detailed information.



Taylosym %24

Oral Solution

Taylosym %24 Oral Solution
240 mg tylosin tartarate

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Taylosym 24% Oral Solution is indicated for the treatment of respiratory infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD) and infectious sinus inflammation in turkeys.

PACKAGING: 500 ml, 1000 ml, 2500 ml, 5000 ml

DOSE: 12,5 ml/ 100 kg total b.w.

WITHDRAWAL TIME: Meat: 2 days (chicken), 5 days (Turkey)
The chicken eggs obtained during the treatment and for a period of 5 days following the last drug administration should not be used for human consumption.

See the leaflet for detailed information.



Tetrasym %80

Water Soluble Powder

Tetrasym %80 Water Soluble Powder
800 mg oxytetracycline

TARGET SPECIES: Calves, Chicken, Turkey, Lamb, Kids

INDICATIONS: TETRASYM 80% Oral Solution Powder is indicated for the treatment of infections caused by sensitive bacteria in calves, chicken, turkey, lamb and kids.

PACKAGING: 500 g and 1000 g

DOSE: 2.5 g/ 100 kg total b.w./ day

WITHDRAWAL TIME: Meat: 7 days.

Egg: zero "0" days.

See the leaflet for detailed information.



Kantel

Solution for Injection

Kantel Solution for Injection
56,80 mg paraziquantel

TARGET SPECIES: Cats, dogs

INDICATIONS: Treatment of cestod infestations in cats and dogs

PACKAGING: 20 ml

DOSE:

Practically, 0.1 ml is injected for 1 kg
body weight. (1 ml/10 kg b.w.)
(s.c.)(i.m.)

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Midokar

Solution for Injection

120 mg imidocarb dipropionate

TARGET SPECIES: cattle, dog, horse

INDICATIONS: It is indicated for the treatment and prevention of babesiosis in cattle, dog and horse.

PACKAGING: 20 ml, 50 ml

DOSE:

Cattle: 1ml/100 kg bw for treatment, 2.5 ml/100 kg bw for prevention (i.m.)(s.c.) Dog: 0.25 ml/10 kg bw for treatment, 0.5 ml/10 kg bw for prevention (i.m.)(s.c.) Horse: 2 ml/100 kg bw for treatment and prevention (i.m.)(s.c.)

WITHDRAWAL TIME: Meat: 213 days, Milk: 21 days

See the leaflet for detailed information.



Klosym

Solution for Injection

Closantel sodium dihydrate equivalent of 50 mg Closantel sodium base

TARGET SPECIES: cattle, sheep

INDICATIONS: Treatment of closantel sensitive trematods, nematods and arthropod larvae invasions.

PACKAGING: 100 ml

DOSE: 1-2 ml/20 kg b.w., s.c. For cattle, max. 20 ml (6 ml in calves) for an injection area. 1-2 ml/20 kg b.w., s.c. for in sheep, max. 5 ml for an injection area.

WITHDRAWAL TIME: Meat: 84 days (cattle), 107 days (sheep).
Not to be used on animals producing milk for human consumption or cattle at last trimester.

See the leaflet for detailed information.



Symec %1

Solution for Injection

Symec %1 Solution for Injection
10 mg Ivermectin

TARGET SPECIES: cattle, sheep

INDICATIONS: It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, subcutaneous nematodes, eye nematode, noca agents, scabies, lice and tick infestations in cattle.

It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, nasal nematodes, scabies and lice infestations in sheep.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Practical dosage is 1 ml for 50 kg bodyweight
Only given subcutaneously.

WITHDRAWAL TIME:Meat: 49 days (cattle), 42 days (sheep) Not to be used within 60 days to birth in animals producing milk intended for human consumption

See the leaflet for detailed information.



Symec Fascio

Solution for Injection

Symec Fascio Solution for Injection
10 mg ivermectin, 100 mg clorsulon

TARGET SPECIES: cattle

INDICATIONS: It is used in the prevention and treatment of gastrointestinal nematodes, lung nematodes (adult and 4th stage larvae), subcutaneous nematode, eye nematode, nocturnal agents (larval stage), lice, scabies and liver trematodes in cattle.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 50 kg/ bodyweight Use subcutaneously only

WITHDRAWAL TIME:

Meat: 66 days,

Not authorised to use on cattle that produce milk for human consumption.

Not to be used within 60 days before giving birth.

See the leaflet for detailed information.



Albesym %10

Oral Suspension

Albesym %10 Oral Suspension
100 mg Albendazole

TARGET SPECIES: cattle, sheep, goat

INDICATIONS: It is indicated to treatment and prevention of infestations caused of gastrointestinal nematode, lungworm and trematodes.

PACKAGING: 1 Liter

DOSE:

Cattle: 7,5 ml/ 100 kg b.w. for gastrointestinal roundworms and lungworms and 10 ml/ 100 kg b.w. for adult flukes

Sheep: 1 ml/ 20 kg b.w. for gastrointestinal roundworms and lungworms and 1,5 ml/ 20 kg b.w. for adult flukes, Dicrocoeliasis dose is 3 ml/20 kg b.w.

Goats, worm and fluke dose is 1,5 ml/ 20 kg b.w. and Dicrocoeliasis dose is 6 ml/40 kg b.w.

WITHDRAWAL TIME:

Meat: 14 days for cattle, 10 days for sheep and goats

Milk: 3 days for cattle, 6 days for sheep and goats (not for human consumption during treatment)

See the leaflet for detailed information.



Albesym 1200

Oral Tablet

Albesym 1200 Oral Tablet
1200 mg Albendazole

TARGET SPECIES: cattle

INDICATIONS: It is indicated to treatment and prevention of infestations caused by worms, liver flukes.

PACKAGING: 10 tablets

DOSE: 1 tablet/ 160 kg b.w. for worms and yung flukes and 1 tablet/ 120 kg b.w.for adult flukes

WITHDRAWAL TIME: Meat: 14 days

Milk: 3 days (6 milkings) (not for human consumption during treatment)

See the leaflet for detailed information.



Medizanid

Oral Tablet

Medizanid Oral Tablet
300 mg Oxfendazole, 600 mg Oxiclosanide

TARGET SPECIES: cattle, sheep

INDICATIONS: Medizanid Oral Tablet is used in prophylaxis and treatment of gastrointestinal and pulmonary infestation by mature and larvae of pinworm, and mature liver flukes, mature *Moniezia* sp. in cattle, camel and sheep.

- a) Liver Flukes
- b) Gastrointestinal pinworms
Stomach
Intestine
Large intestine
- c) Pulmonary pinworms
- d) Tapeworm

PACKAGING: 10, 20, 30, 50 tablet

DOSE:

Lamb: ½ tablet for 25 kg

Sheep: 1 tablet for 50 kg

Calf-Steer: 2 tablets for 100 kg

Cattle: 10 tablet for 500 kg

WITHDRAWAL TIME:Meat: 28 days (cattle) , 21 days (sheep)

Milk: 5 days (cattle) Not to be used for dairy cows producing milk for human consumption.

See the leaflet for detailed information.



Wormkill

Oral Tablet

Wormkill Oral Tablet

50 mg Praziquantel, 144 mg Pirantel pamoat 150 mg Febantel

TARGET SPECIES: dog

INDICATIONS: WORMKILL Oral Tablet is indicated for the control of and fighting against the parasitic invasions caused by the following nematode and cestode species in dogs. Ascarid: *Toxocara canis*, *Toxascaris leonina* (mature and immature) Hookworm: *Uncinaria stenocephala*, *Ancylostoma caninum* (mature) Whipworm: *Trichuris vulpis* (mature) Cestode: *Echinococcus granulosus*, *Echinococcus multilocularis*, *Dipylidium caninum*, *Taenia* sp.

PACKAGING: 50 tablets

DOSE: Practical dosage for dogs: 1 tablet for 10 kg bodyweight. Tablets may also be administered half according to the weight of the animals.

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Flumetsym

Pour on Solution

Flumetsym Pour on Solution
10 mg flumethrin

TARGET SPECIES: cattle, sheep

INDICATIONS: Flumetsym Pour on Solution is used for the control and treatment against biting flies, chewing and biting lice (Haematopinus eury-sternus, Linognathus vituli, Bovicola bovis, B. ovis, Melophagus ovinus), ticks, mange mites (scabies) and other ecto parasites related infections in cattle, sheep and dogs.

PACKAGING: 100 ml, 500 ml

DOSE: The application dose is 1 mg/kg b.w. for tick control in cattle and sheep, and 2 mg/kg body weight for scabies and lice control.

WITHDRAWAL TIME:

Meat: 21 days (sheep) 5 days (cattle)

Milk: 8 days

Not authorised to use on sheep that produce milk for human consumption.

See the leaflet for detailed information.



Amprolsym %25

Oral Solution

Amprolsym %25 Oral Solution
250 mg Amprolium

TARGET SPECIES: Chicken


INDICATIONS: Treatment and prevention of coccidiosis caused by *Eimeria tenella*, *E. Necatrix*, *E. Acervulina*, *E. Maxima*, *E. Brunetti* in chicken

PACKAGING: 100 ml, 500 ml, 1000 ml and 5000 ml

DOSE: 300 ml/1000 kg total b.w. for 7 days

WITHDRAWAL TIME:
"0" days for milk and egg

See the leaflet for detailed information.



Symcox

Oral Solution

Symcox Oral Solution
25 mg Toltrazuril

TARGET SPECIES: chicken, turkey

INDICATIONS: It is indicated for the treatment of and protection against coccidiosis in chickens and turkeys. Below are the most important coccidiosis agents, which are included in the spectrum of action of Symcox Oral Solution.; For chickens: Eimeria acervulina, Eimeria brunetti, Eimeria maxima, Eimeria mivati, Eimeria necatrix, Eimeria tenella For turkeys: Eimeria adenoides, Eimeria meleagrimitis

PACKAGING: 100 ml, 1000 ml

DOSE: 280 ml product/1000 kg total body weight

WITHDRAWAL TIME:

Meat: 21 days

It should be administered to chickens and turkeys producing egg for human consumption.

See the leaflet for detailed information.



IPM Ksilazin %2

Solution for Injection
Veterinary Sedative, Hypnotic

IPM Ksilazin Solution for Injection
20 mg xylazine

TARGET SPECIES: Horse, Cattle, Dog, Cat

INDICATIONS: For sedation, muscle relaxation and analgesia in minor operations, as well as premedication for general anesthesia

PACKAGING: 50 ml

DOSE: It is administered by intravenous, intramuscular and subcutaneous routes in target species.
Please read leaflet for detailed information

WITHDRAWAL TIME:Meat: 1 day
Milk "0" day

See the leaflet for detailed information.



Furosym

Solution for Injection

Furosym Solution for Injection
20 mg furosemide

TARGET SPECIES: cattle, horse, cat, dog

INDICATIONS: Furosym Solution for Injection is used in order for elimination of fluids, accumulating and causing the formation of edema due to the disorders in circulatory system, liver, kidneys and any other reasons, from the tissues, body cavities, joints, tendons and surfaces through the diuresis, and in order for acceleration of diuresis in case of intoxications, and also it is used in case of emergencies such as acute lung and brain edema.

PACKAGING: 20 ml, 50 ml

DOSE: Practical dosage: 2,5 - 5 ml/100 kg/bw for cattle and horses (slowly intravenous), 1,25 -2,5 ml/10 kg/bw for dog and cats (slowly intravenous or intramusculary)

WITHDRAWAL TIME: "0 days" meat and milk (i.v.)

See the leaflet for detailed information.



Symatro

Solution for Injection

Symatro Solution for Injection
2 mg Atropine

TARGET SPECIES: Horse, cat, dog

INDICATIONS: For stimulation of hearth movements in horse, dog and cats,
at cases of pain, bladder, uretra and kidney spasms colics and constrictions as spasmolitic,
at organic phosphorus poisoning with aracoline, pilocarpin and neostigmin poisonings as antidote
at general anesthesia for reduction of saliva, for prevention of vago-vagal reflexes and vagal tonus negative effects on hearth as preanesthetic medication.

PACKAGING: 20 ml and 50 ml

DOSE:

Horse 1,5 - 3 ml/100 kg l.w. 30-60 µg/kg b.w.
Dog 0,15 - 0,25 ml/10 kg l.w. 30-50 µg/kg b.w.
Cat 0,06- 0,1 ml/ 4 kg l.w. 30-50 µg/kg b.w.

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Symbuton

Solution for Injection

100 mg/ml menbuton

TARGET SPECIES: cattle, horse, sheep, goat

INDICATIONS: Stimulation of hepato-digestive activity in case of digestive disorders and hepatic insufficiency

PACKAGING: 100 ml

DOSE:

Cattle : 1 ml for 15-20 kg/ bodyweight by intravenously
 Sheep, goat and calves : 1 ml for 10 mg/kg bodyweight by deep intramuscular injection or intravenous slowly
 Horse : 1 ml for 20-40 kg/ bodyweight by intravenous injection

WITHDRAWAL TIME: '0' days for meat and milk

See the leaflet for detailed information.



Adecure AD₃E

Solution for Injection

Adecure AD3E Solution for Injection
500.000 IU Vitamin A, 75.000 IU Vitamin D3, 50 mg Vitamin E

TARGET SPECIES: cattle, sheep, goat, rabbit

INDICATIONS: A, D3 and E avitaminosis and treatment for related diseases and malnutritions.

PACKAGING: 100 ml

DOSE:

Calves: 1-3 ml

Cattle: 5-8 ml

Lamb - Goat: 0,25 - 0,5 ml

Sheep - Goat: 1 ml

Rabbit: 0,1 - 0,25 ml

WITHDRAWAL TIME:

Zero "0" day for meat and milk

See the leaflet for detailed information.



Adezek

Solution for Injection
Vitamin

Adezek Solution for Injection

80 mg Vitamin A (200.000 IU), 0,7875 mg Vitamin D3 (30.000 IU),
20 mg Vitamin E

TARGET SPECIES: cattle, sheep, goat, rabbit

INDICATIONS:Adezek Solution for Injection is indicated for to prevent deficiencies and treatment of diseases due to avitaminosis of these vitamins in cattle, sheep, goat and rabbits.

PACKAGING:100 ml

DOSE:

Cattle: 5-10 ml

Calves: 2-4 ml

Sheep- Goat: 1-2 ml

Lamb-rabbit: 0.5 ml

WITHDRAWAL TIME:

28 days for meat

5 days for milk

See the leaflet for detailed information.



Bevitbar

Solution for Injection

Bevitbar Solution for Injection

100 mg Vitamin B1, 10 mg Vitamin B6

TARGET SPECIES:cattle, sheep, goat, cat, dog

INDICATIONS:Bevitbar Solution for Injection is used against Vitamin B1 (Thiamin HCl) and Vitamin B6 (Pyridoxine HCl) deficiencies or as supportive to treatment in cattle, sheep, goat, cat and dog.

PACKAGING:100 ml

DOSE:

Cattle - Horse 400 kg 10 -20 ml daily

Steer - Heifer 200 kg 5 - 10 ml daily

Sheep - Goats 100 kg 2,5 - 5 ml daily

Cat - Dog 5 kg 1 -2 ml daily

WITHDRAWAL TIME:"0" days for milk and meat

See the leaflet for detailed information.



Symvit-C

Solution for Injection

Symvit-C Solution for Injection
200 mg Vitamin C (Ascorbic acid)

TARGET SPECIES: cattle, sheep, horse, goat, cat, dog

INDICATIONS: It is used to treatment and prevention diseases caused by ascorbic acid (vitamin C) deficiency in animals. It is used in clinical and supportive treatment in cases of developmental and adaptation deficiencies, anemia, hypocalcemia, mycotoxiosis and intoxication

PACKAGING: 50 ml, 100 ml

DOSE:

Cattle, 7 - 12 ml /day i.m , s.c , i.v twice a week during 3-6 weeks

Steer, 2 - 4 ml/day i.m, s.c, i.v twice a week during 3 weeks

Sheep, 2 - 4 ml/day i.m , s.c twice a week during 3 weeks

Goat, 2 - 4 ml/day i.m , s.c twice a week during 3 weeks

Horse, 10 -20 ml /day i.m, i.v

Cat, 0,3 - 0,6 ml/day i.m, i.v

Dog, 0,3 – 2,5 ml/day i.m, i.v

WITHDRAWAL TIME: "0" days for meat and milk

See the leaflet for detailed information.



Symfos

Solution for Injection

Symfos Solution for Injection
100 mg Butaphosphan, 0.05 mg Vitamin B12 (Cyanocobalamine)

TARGET SPECIES: cattle, horse, cat, dog

INDICATIONS: It is used as a supplement in muscle, metabolism or reproductive system disorders where the use of phosphorus and cyanobalamin is required in cattle, horse, cats and dogs.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 2-5 ml/ 100 kg b.w. (i.v.) in horses and cattle.
0,35-0,6 ml/10 kg b.w. in foals and calves.
0,25-2,5 ml/ 10 kg b.w. (i.v., i.m., s.c.) in dogs.
0,5-5 ml/ 5 kg b.w. (i.v., i.m., s.c.) in cats.
Doses must be split in half with 1-2 weeks delay for chronic conditions in cats and dogs.

WITHDRAWAL TIME: '0' days for meat and milk (cattle, horse)

See the leaflet for detailed information.



Symcal

Solution for Injection

Symcal Solution for Injection

190 mg calcium gluconate, 45 mg calcium glucoheptonate, 60 mg magnesium chloride, 4 mg butaphosphane, 10 mg calcium d-saccharide

TARGET SPECIES: Cattle, Horse, Calves, Foals, Sheep, Goat, Dog

INDICATIONS: It is indicated for the protection from and treatment of acute and chronic metabolism disorders resulted from the deficiency of calcium, magnesium and phosphor in cattle, horses, sheep, goats and dogs.

PACKAGING: 100 ml, 250 ml

DOSE: Use intramuscularly, intravenously or subcutaneously
Practical dosage: 1 ml/1 kg bw

WITHDRAWAL TIME: "0" days for meat and milk (cattle, sheep, goat)

See the leaflet for detailed information.



Badeljin

Solution for Injection

Badeljin Solution for Injection
500 mg metamisole sodium

TARGET SPECIES: cattle, horse, dog

INDICATIONS: Badeljin Solution for Injection is used in the symptomatic treatment of the following cases in cattle, horses and dogs.

As analgesic, Spasmolithically, Antipyretically, Antirheumatic

PACKAGING: 50 ml and 100 ml

DOSE:

For horses (per 100 kg l.w.) 4 -10 ml For cattle (per 100 kg l.w.) 8 ml
Foal -bovine (per 50 kg l.w.) 2-4 ml For dogs (per 10 kg l.w.) 1 ml

WITHDRAWAL TIME: Meat: 12 days Do not use in lactating cows producing milk for human consumption

See the leaflet for detailed information.



Ketozek

Ketozek Solution for Injection
100 mg ketoprofen

TARGET SPECIES: cattle, horse

INDICATIONS: Diseases affecting the osteoarticular and muscular-skeletal system associated with acute pain and inflammation, after calving.

PACKAGING: 100 ml

Solution for Injection
Non-steroidal Anti-inflammatory, Antiromatizmal

DOSE:

For Cattle 3 ml/ 100 kg b.w. use intramuscular or intravenously, for up to 3 days For horse 1 ml/ 45 kg b.w. for horse use intravenously, for 3-5 days.

WITHDRAWAL TIME: Meat: 4 days Milk: Zero days.

See the leaflet for detailed information.



Melosym

Solution for Injection

Melosym Solution for Injection
5 mg meloxicam

TARGET SPECIES: cattle, dog, cat

INDICATIONS: In cattle; It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory tract infection. In dogs; It is used to relieve inflammation and relieve pain in acute and chronic musculoskeletal diseases, and to reduce inflammation and pain after orthopedic and soft tissue operations. in cats; It is used to reduce post-operative pain after ovariohysterectomy and mild soft tissue operations.

PACKAGING: 20 ml, 50 ml, 100 ml

DOSE: Cattle: 10 ml/100 kg bw subcutaneously or intravenously
Dogs: 04 ml/10 kg bw subcutaneously
Cat: 0.12ml/2 kg bw subcutaneously

WITHDRAWAL TIME:

Meat: 15 days (Cattle) **Milk:** 5 days (10 milkings) (cattle)

See the leaflet for detailed information.



Melosym X8

Solution for Injection

Melosym x8 Solution for Injection
40 mg Meloxicam

TARGET SPECIES: cattle, horse

INDICATIONS: It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory infection.

PACKAGING: 100 ml

DOSE: Cattle: 1.25 ml / 100 kg b.w. by single dose subcutaneous (sc) or intravenous (iv) injection.

Horse: It is administered by a single dose intravenous (iv) i 1.5 ml / 100 kg b.w.

WITHDRAWAL TIME:

Meat: 15 days (Cattle) **Milk:** 5days (10 milkings) (cattle)

See the leaflet for detailed information.



Symful

Solution for Injection

Symful Solution for Injection
50 mg flunixin

TARGET SPECIES: cattle, horse

INDICATIONS: Horse: Painful and inflammatory conditions of the skeletal-muscular system and colic.

Cattle: Accompanying proper treatment, to reduce clinical symptoms formed during pulmonary infections.

PACKAGING: 100 ml, 250 ml

DOSE: Intravenous application should be slow and administered at body temperature.

2 ml/ 50 kg b.w. (slow i.v., i.m.) for 1-3 days in cattle.

1 ml/ 50 kg b.w. (slow i.v.) for 1-5 days in horses.

WITHDRAWAL TIME:Meat: 31 days (i.m.), 10 days (i.v.)

Milk: 36 hours (i.m.), 24 hours (i.v.)

See the leaflet for detailed information.







İTOSB (İstanbul Tuzla Organize Sanayi Bölgesi) 13. Cadde No:1/B,
34959 Tepeören, Tuzla, İstanbul - TÜRKİYE

T: +90 0216 290 22 44 info@ipmilac.com

www.ipmilac.com

