

ANIMAL HEALTH
PRODUCTS



ipm



Index

ANTIBACTERIAL

Injectable Products

Ceftiforce Suspension for Injection	6
Flormeg Solution for Injection.....	7
Florsym Solution for Injection.....	8
Linkosym Solution for Injection	9
Marboforce Solution for Injection	10
Oksipirimavet LA Solution for Injection.....	11
Pentril %10 Solution for Injection	12
Ribogen Solution for Injection.....	13
Spiraforce Solution for Injection.....	14
Sulfakill Solution for Injection	15
Symtil 300 Solution for Injection.....	16
Taylobar Solution for Injection	17
Tulasym Solution for Injection	18

Antibacterial Intramammary

Ampriforce DC Intramammary Suspension	19
---	----

Oral Products

Doksisym %50 Water Soluble Powder.....	20
Doksisym Oral Tablet.....	21
Doksol %10 Oral Solution	22
Enrosym %10 Oral Solution.....	23
Enrosym %20 Oral Solution.....	24
Enrosym %80 Water Soluble Powder	25
Enrosym Oral Tablet	26
Erizol Water Soluble Powder.....	27
Florsym %30 Oral Solution.....	28
Kolisym %24 Water Soluble Powder	29
Linkosym Water Soluble Powder	30
Metrisym Oral Tablet	31
Neosym %50 Water Soluble Powder.....	32
Paripto Powder for Oral Solution	33
Spiramezol Oral Tablet	34
Sulfasym Oral Suspension.....	35
Symtay Water Soluble Powder.....	36
Symtil %30 Oral Solution.....	37
Taylosym %24 Oral Solution	38
Tetrasym %80 Water Soluble Powder	39

ANTIPARASITIC

Injectable Products

Kantel Solution for Injection	40
Midokar Solution for Injection	41
Klosym Solution for Injection.....	42
Symec %1 Solution for Injection	43
Symec Fascio Solution for Injection	44

Oral Products

Albesym %10 Oral Suspension	45
Albesym 1200 Oral Tablet.....	46
Levasym Oral Suspension.....	47
Medizanid Oral Tablet	48
Prazifen KK Oral Tablet	49
Wormkill Oral Tablet.....	50

Topical Solutions

Eprinosym Pour-On Solution	51
Flumetsym Pour-on Solution	52

ANTICOCCIDIAL

Amprolsym %25 Oral Solution.....	53
Symcox Oral Solution	54

CLINICAL MEDICATIONS

IPM Ksilazin Solution for Injection.....	55
Furosym Solution for Injection.....	56
Symatro Solution for Injection	57
Symbuton Solution for Injection	58

VITAMIN

Adecure AD3E Solution for Injection.....	60
Adezek Solution for Injection	61
Bevitbar Solution for Injection	62
Selesym Solution for Injection.....	63
Symvit-C Solution for Injection	64
Symfos Solution for Injection.....	65

MINERAL

Symcal Solution for Injection	66
-------------------------------------	----

ANALGESIC- ANTI-INFLAMMATORY

Badeljin Solution for Injection	67
Ketozek Solution for Injection.....	68
Melosym Solution for Injection	69
Melosym X8 Solution for Injection.....	70
Symful Solution for Injection.....	71



ANIMAL HEALTH PRODUCTS

ipm

Ceftiforce

Suspension for Injection

50 mg/ml Ceftiofur HCl

TARGET SPECIES: Cattle

INDICATIONS: It is used in cattle for the treatment of respiratory and soft tissue infections caused by ceftiofur-sensitive bacteria. It is used in the treatment of Mannheimia sp. (Pasteurella haemolytica), Pasteurella multocida, Respiratory tract infections caused by Haemophilus somnus, Acute interdigital necrobacillosis (fork rot, pododermatitis) caused by Fusobacterium necrophorum and Bacterioides melaninogenicus, acute mastitis with postnatal e coli, Arcanobacterium necropoygenium, and Arcanobacterium bacterium acute puerperal metritis.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml/50 kg/bw intramuscularly or subcutaneously, one application for every 3-5 days

WITHDRAWAL TIME:

Meat: 8 days

Milk: 0 day



Flormeg

Solution for Injection

300 mg/ml Florfenicol 16.5 mg/ml Flunixin

TARGET SPECIES: Cattle

INDICATIONS: It is used in cattle for the treatment of respiratory infections caused by Mannheimia haemolytica, Pasteurella multocida, Mycoplasma bovis and Histophilus somni.

PACKAGING: 100 ml, 250 ml

DOSE: 2 ml/15 kg bw/day, one application subcutaneously

WITHDRAWAL TIME: Meat: 46 days.

Milk: Not authorized for use in lactating animals producing milk for human consumption.

See the leaflet for detailed information.



Florsym

Solution for Injection

300 mg/ml Florfenicol

TARGET SPECIES: Cattle, Sheep

INDICATIONS:

Cattle: Treatment and metaphylaxis of respiratory tract infections caused by florfenicol-susceptible strains of Mannheimia haemolytica, Pasteurella multocida and Histophilus somni. Before metaphylactic use, the presence of disease in the flock should be ensured.

Sheep: respiratory tract infections from Mannheimia haemolytica and Pasteurella multocida.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml/ 15 kg b.w. (i.m.),
2 ml/ 15 kg b.w.(s.c. for cattle only) treatment should continue at 48 hours intervals.

Sheep: 1 ml/15 kg bw (i.m.) for 3 days

WITHDRAWAL TIME:Meat: 30 days (i.m., cattle), 39 days (sheep, 44 days (s.c., cattle) Do not use in lactating cattle producing milk (including dry period) for human consumption

See the leaflet for detailed information.



Linkosym

Solution for Injection

Lincomycin hydrochloride (equivalent to 50 mg/ml lincomycin base)
Spectinomycin sulfate tetrahydrate (equivalent to 100 mg/ml spectinomycin base)

TARGET SPECIES: Sheep, Goat, Calves, Cattle, Dog, Cat, Broiler

INDICATIONS: It is used in the treatment of respiratory, digestive, urinary system infections and soft tissue infections and arthritis caused by bacteria sensitive to lincomycin and spectinomycin in target species.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE:

Sheep-Goat: 1 ml for 10 kg bodyweight during 3-5 days

Calf Cattle: 1 ml for 10 kg bodyweight, intramuscularly, daily during 3-5 days (first day 2 injections)

Dog-Cat: 1 ml for 5 kg body weight 1-2 a day, during 3-7 days.

Chickens and Turkey: 1 ml for 5 kg, subcutaneously for 3 days

WITHDRAWAL TIME:

Meat: 21 days for Cattle 14 days for Sheep, Goat, Broiler

Not to be used in sheep producing milk intended for human consumption

Not to be used in chicken producing egg intended for human consumption

See the leaflet for detailed information.



Marboforce

Solution for Injection

100 mg/ml Marboploxacin

TARGET SPECIES: Cattle

INDICATIONS: It is indicated for respiratory tract infections caused by *Pasteurella multocida*, *Mannheimia haemolytica* ve *Histophilus somni* and used for the treatment of acute mastitis and infections caused by *Escherichia coli* and *Mycoplasma bovis*.

PACKAGING: 100 ml

DOSE: For respiratory track infections dose is 2 ml/25 kg b.w. (i.m.). For *Mycoplasma bovis* infections and acute mastitis dose is 1 ml/50kg b.w./ day (i.m.,s.c.), should be applied for 3 days.

WITHDRAWAL TIME:

Meat: 3 days (for 8 mg/ kg b.w. treatment), 6 days (for 2 mg/ kg b.w. treatment)

Milk: 3 days (6 milkings, 8 mg/ kg b.w.), 1,5 days (3 milkings, 2 mg/ kg b.w.)

See the leaflet for detailed information.



Oksipirimavet LA

Solution for Injection

200 mg/ml Oxytetracycline

TARGET SPECIES: Cattle, Sheep, Goat

INDICATIONS: It indicated to treatment of systemic, regional and respiratory system infections caused by bacteria sensitive to oxytetracycline.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 10 kg/bw (i.m)

WITHDRAWAL TIME:

Meat: 28 days

Milk: 10 days (cattle), 7 days (sheep, goat)

See the leaflet for detailed information.



Pentril %10

Solution for Injection

100 mg/ml Enrofloxacin

TARGET SPECIES: Cattle, Sheep, Goat

INDICATIONS: It is used for the treatment of sensitive gram negative, gram positive and Mycoplasma related infections in cattle and sheep and goats.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Practical dose: 1 ml for 20 kg bw/day Product may given subcutaneously or intravenously. Treatment should continue for 3-5 days.

WITHDRAWAL TIME:

Meat: 5 days (cattle i.v.), 12 days (cattle s.c.), 4 days (sheep), 6 days (goats)

Milk: 3 days (sheep, cattle i.v.), 4 days (goats, cattle s.c.)

See the leaflet for detailed information.



Ribogen

Solution for Injection

100 mg/ml Gentamycine

TARGET SPECIES: Cattle, Horse, Dog, Cat

INDICATIONS: It is used against gentamicin sensitive bacteria causing respiratory, digestive and urogenital system infections, septicaemia and other soft tissue infections treatment in cattle, horse, dog and cat.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Cattle: 2 ml / 100kg body weight for 3-7 days

Horse: 6.5 ml/100kg bw for 3-5 days

Dog and cat: 0.2ml / 5 kg body weight for 3-5 days

WITHDRAWAL TIME: Meat: 139 days(cattle), 103 days(calves) for 3 injections,

192 days for 6 injections, 214 days for 7 injections

Milk: 7 days

See the leaflet for detailed information.



Spiraforce

Solution for Injection

600,000 IU/ml Spiramycin

TARGET SPECIES: Cattle

INDICATIONS: In lactating cows, it is used to treat acute clinical mastitis caused by strains of *Staphylococcus aureus* susceptible to Spiramycin, and respiratory tract infections caused by *Pasteurella multocida* and *Mannheimia haemolytica*.

PACKAGING: 100 ml, 250 ml

DOSE: When used at 5 ml / 100 kg b.w., twice with a 24-hour interval or 5 ml / 30 kg b.w., twice with a 48-hour interval:

WITHDRAWAL TIME:

Meat: 75 days

Milk: 13.5 days (27 milkings)

See the leaflet for detailed information.



Sulfakill

Solution for Injection

215.8 mg/ml Sulfadimidin Sodium , 40 mg/ml Trimetophrim

TARGET SPECIES: Cattle, Horse, Cat, Dog

INDICATIONS: Sulfakill Solution for Injection is indicated for the infections caused by susceptible microorganisms

PACKAGING: 50 ml, 100 ml ve 250 ml

DOSE: 1 ml/10-15 kg b.w. for cattle and horses; 0,2 ml/3 kg bw for cats and dogs.

WITHDRAWAL TIME: Meat: 12 days **Milk:** 5 days

See the leaflet for detailed information.



Symtil 300

Solution for Injection

300 mg/ml Tilmycosin

TARGET SPECIES: Cattle, Sheep

INDICATIONS: SYMTİL 300 Solution for injection is indicated for treatment of respiratory disease (BRD) caused by bacteria susceptible to tilmycosin, treatment of respiratory disease associated with Pasteurella haemolytica, treatment of interdigital necrobacillosis and foot rot caused by Fusobacterium necrophorus, prevention of abortus caused by Chlamidia psittaci and as parenteral support in the treatment of mastitis and metritis in cattle and sheep.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 30 kg b.w.

WITHDRAWAL TIME:

Meat: 70 days (cattle), 42 days (sheep)

Milk: 18 days (sheep) 36 days(cattle)

See the leaflet for detailed information.



Taylobar

Solution for Injection

200 mg/ml Tylosin

TARGET SPECIES: Cattle, Sheep, Goat, Cat, Dog

INDICATIONS:

Cattle (adult): metritis caused by gram positive bacteria, respiratory system infections, Streptococcus spp., Staphylococcus spp. Treatment of mastitis caused by Fusobacterium necrophorum and interdigital necrobacillosis caused by Fusobacterium necrophorum, **Calf:** Necrobacillosis (Fusobacterium necrophorum) and respiratory system infections

Sheep-goat: metritis caused by gram positive bacteria, respiratory system infections, Gram positive bacteria or Mycoplasma spp. mastitis caused by Dog: Respiratory tract infections and otitis caused by susceptible bacteria Cat: respiratory tract infections caused by susceptible bacteria

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE:

Applied intramuscularly and only applied slow intravenous for cattle.

Cattle: 2.5-5 ml/100 kg bw

Sheep-goat: 2.5 ml/50 kg bw

Cat-dog: 0.35-0.55 ml/10 kg bw

WITHDRAWAL TIME:Meat: 28 days (cattle) 42 days (sheep, goat)

Milk: 108 hours

See the leaflet for detailed information.



Tulasym

Solution for Injection

100 mg/ml Tulatromycin

TARGET SPECIES: Cattle, Sheep

INDICATIONS:

Cattle; It is indicated for the treatment and metaphylaxis of bovine respiratory disease (BRD) caused by Mannheimia haemolytica, Pasteurella multocida, Histophilus somni and Mycoplasma bovis. It is indicated for the treatment of infectious keratoconjunctivitis caused by Morexella bovis sensitive to tulatromycin

Sheep; It is indicated for the early treatment of pododermatitis caused by virulent Dichelobacter nodosus

PACKAGING: 100 ml, 250 ml

DOSE: Cattle: 2,5 mg tulatromycin/kg b.w. (1ml/40 kg b.w.) single dose via s.c.

Sheep: 2,5 mg tulatromycin/kg b.w. (1ml/40 kg b.w.) single dose via i.m. route

WITHDRAWAL TIME:

Meat: Cattle: 22 days, **Sheep:** 16 days

Not to be used in animals producing milk intended for human consumption or animals expected to give birth in two months that will produce milk for human consumption.

See the leaflet for detailed information.



Ampiforce DC

Intramammary Suspension

Cloxacillin Benzathine Equivalent to 600 mg Cloxacillin,
Ampicillin Trihydrate Equivalent to 300 mg Ampicillin

TARGET SPECIES: It is used in cows in the dry period.

INDICATIONS: It is used immediately after the last milking of a lactation to treat existing mastitis and to protect against further infections during the dry period. It is active against Gram-positive and Gram-negative organisms associated with mastitis, and is effective against *Streptococcus agalactiae* and other *Streptococcus* species, penicillin-resistant and susceptible *Staphylococci*, *Arcanobacterium* spp., *Escherichia coli*, and other susceptible Gram-negative bacteria.

PACKAGING: In cardboard packages of 1, 4 and 50 pieces or in polypropylene buckets of 48 and 120 pieces.

DOSE: It is administered as an intraudder infusion to heifers and lactating cows. It should be administered as a tube into each udder after the final milking, after the teat has been cleaned and disinfected. The animal is properly restrained. The teats are cleaned and disinfected. The tip of the syringe is inserted into the teat opening, BUT NOT INSERTION INTO IT. When the plunger is pushed, the product reaches the udder tissue through the teat.

WITHDRAWAL TIME:

Dry period short and cows <49 days before calving: not for use.

Milk: Calving ≥ 49 days: 156 hours

Calving <49 days: 49 days + 156 hours from treatment

Accidental use during lactation: milk must not be used for at least 28 days and until residue levels fall below the limits.

Meat: 28 days

See the leaflet for detailed information.



Doksisym %50

Powder for Oral Solution

577 mg/g Doxycycline hyclate (equivalent to 500 mg/g base)

TARGET SPECIES: Non-Ruminating Calves, Chicken, Turkey

INDICATIONS: It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

PACKAGING: 1 kg

DOSE:

Calves; 17.3 mg product/b.w/day via oral route during 3 or 5 days

Chicken; 43.3 mg product/b.w/day via oral route during 3 or 5 days

Turkey; 50.2 mg product/b.w/day via oral route during 3 or 5 days

WITHDRAWAL TIME:

Meat: 7 days (calves), 5 days (chicken), 12 days (turkey)

Egg: Do not use in animals producing egg for human consumption.

Milk: Do not use in animals producing milk for human consumption.
Not to be used within 4 weeks before egg production period.

See the leaflet for detailed information.



Doksisym

Oral Tablet

115,388 mg/tb Doxycycline Hyclate

TARGET SPECIES: Dogs

INDICATIONS: It is indicated for treatment of respiratory system infections such as rhinitis, tonsillitis caused by the doxycycline-sensitive microorganisms

PACKAGING: 10 tablets (1x10) or 30 tablets (3x10) blister packaging in cardboard box

DOSE:

1 tablet/10 kg bw for 5 days

WITHDRAWAL TIME:

Not applicable

See the leaflet for detailed information.



Doksol %10

Oral Solution

100 mg/ml Doxycycline

TARGET SPECIES: Non-Ruminating Calves, Chicken, Turkey

INDICATIONS: It is indicated to treatment of respiratory and digestive system infections on calves, broilers and turkeys

PACKAGING: 100 ml, 1 L

DOSE:

Recommended dose of product is mixed in water

Calves; 0.08 ml product/b.w/day via oral route during 3-5 days

Chicken; 0.21 ml product/b.w/day via oral route during 3-5 days

Turkey; 0.25 ml product/b.w/day via oral route during 3-5 days

WITHDRAWAL TIME:

Meat: 7 days (calves), 5 days (chicken), 12 days (turkey)

Egg: Do not use in animals producing egg for human consumption.

Milk: Do not use in animals producing milk for human consumption.

Not to be used within 4 weeks before egg production period.

See the leaflet for detailed information.



Enrosym %10

Oral Solution

100 mg/ml Enrofloxacin

TARGET SPECIES: Chicken, Turkey, Rabbit


INDICATIONS: It is indicated for the treatment of the infections caused by gram negative and positive bacteria and Mycoplasma spp. susceptible to enrofloxacin in broiler turkeys, chickens and rabbits.

PACKAGING: 100 ml, 2,5L , 3L

DOSE: The daily dose to be administered is 10 mg/kg body weight by adding drinking water for 3-5 days (5 days for rabbit).

WITHDRAWAL TIME:Meat: 7 days (chicken), 13 days (turkey), 3 days (rabbit) Should not be given to chickens and turkeys producing eggs for human consumption, not to be used within 14 days to egg laying period.

See the leaflet for detailed information.



Enrosym %20

Oral Solution

200 mg/ ml Enrofloxacin

TARGET SPECIES: Chicken, Turkey, Rabbit

INDICATIONS: It is used in the treatment of respiratory and gastro intestinal system infections caused by susceptible bacteria in chicken, turkey and rabbits.

PACKAGING: 100 ml ,3 L

DOSE:


Chicken and Turkey: 10 mg/kg body weight, mixed into drinking water.

Rabbit: 10 mg/kg body weight daily, for 5 consecutive days.

WITHDRAWAL TIME: Meat: 7 days (chicken), 13 days (turkey), 3 days (rabbit)

Egg: Do not use in animals producing egg for human consumption. Not to be used within 14 days to egg laying period.

See the leaflet for detailed information.



Enrosym %80

Powder for Oral Solution

800 mg/g Enrofloxacin

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Enrosym 80% Powder for Oral Solution is indicated for the treatment of infections caused by gram-negative and gram-positive bacteria that are enrofloxacin-sensitive, infections caused by mycoplasmas and secondary bacterial infections accompanying viral infections in broiler chickens and broiler turkeys. It is indicated mainly for the following: CRD (Chronic Respiratory Disease), mycoplasmosis, coliseptisemia, colibacillosis, pasteurellosis

PACKAGING: 1 kg

Dose: 1,25 g/100 kg total b.w./day

WITHDRAWAL TIME: Meat: 12 days (chicken), 14 days (turkey).

Not to be used in chicken and turkey producing egg intended for human consumption

See the leaflet for detailed information.



Enrosym

Oral Tablet

150 mg/tb Enrofloxacin

TARGET SPECIES: Dogs

INDICATIONS:

It is indicated to treat following infections.

In dogs; single or combined respiratory system, lower urinary tract (single or with prostatitis), upper urinary tract infections caused by Escherichia Coli or Proteus mirabilis, skin infections, superficial and deep pyodermas, otitis externa caused by gram positive and gram negative bacteria .

PACKAGINGS:

10 tablets (1x10) or 30 tablets (3x10)
blister packaging in cardboard box

DOSE:

1 tablet/30 kg bw for 10-21 days
depending on the severity and type of infection.

WITHDRAWAL TIME:

Not applicable

See the leaflet for detailed information.



Erizol

Powder for Oral Solution

509 mg/ g Eritromycin

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Treatment of infections associated with Mycoplasma

PACKAGING: 100 g and 1 kg

DOSE: It is used as 20-40 mg/g erythromycine base/day via oral route during 3 or 5 days
Practically 39-79 g/1000 kg b.w.

WITHDRAWAL TIME:

Meat: 3 days (chicken), 6 days (turkey)

Egg: 5 days

See the leaflet for detailed information.



Florsym %30

Oral Solution

300 mg/ml Florfenicol

TARGET SPECIES: Chicken

INDICATIONS: Florsym 30% Oral Solution is indicated for the treatment of respiratory diseases, sacculitis and colisepticemia caused by E. Coli.

PACKAGING: 1000 ml and 2500 ml

DOSE: 100ml/1500 kg total b.w./ day

WITHDRAWAL TIME:

Meat: 5 days

Not to be used in chicken producing egg intended for human consumption

See the leaflet for detailed information.



Kolisym %24

Powder for Oral Solution

4.800.000 IU /g Kolistin

TARGET SPECIES: Chicken, Turkey, Calves, Lambs

INDICATIONS: It is indicated for the treatment of digestive system infections caused by Gram negative bacteria including especially E. Coli and Salmonella species in chicken, turkey, calves and lambs.

PACKAGING: 1 kg

DOSE:

15,625g/1000 kg total b.w. for chicken and turkey 3-5 days

1 g/48 kg b.w. is sufficient in calves and lambs

WITHDRAWAL TIME:

Meat: 1 day

Egg: zero "0" days

not to be used on animals that produce milk for human consumption.

See the leaflet for detailed information.



Linkosym

Powder for Oral Solution

222 mg/ ml Lincomycin, 444 mg/ml Spectinomycin

TARGET SPECIES: Chicken

INDICATIONS: Linkosym Powder for Oral Solution is indicated for the treatment of the infections caused by sensitive bacteria, chronic respiratory infections (CRD) caused especially by Mycoplasma sp., and CRD complex caused both by Mycoplasma sp. and coliform microorganisms, avian cholera and other coliform infections in broiler chickens and poults, as well as air sac infections in turkeys.

PACKAGING: 150 g

DOSE: 7.5 g /100 kg total b.w. for chicken

WITHDRAWAL TIME:

Meat: 5 days

Not to be used in poultry producing egg intended for human consumption

See the leaflet for detailed information.



Metrisym

Oral Tablet

Metrisym Oral Tablet
1000 mg/ tb Sulfadiazin 200 mg/tb Trimethoprim

TARGET SPECIES: Calves

INDICATIONS: Metrisym Oral Tablet is mainly indicated for the treatment of bacterial diarrhea in calves, it can also used in the treatment of acute salmonellosis and bacterial pneumonia.

PACKAGING: 2x 5 tablets

DOSE:

1 tablet/ 40 kg b.w./ day
during 5 days

WITHDRAWAL TIME: Meat: 15 days

See the leaflet for detailed information.



Neosym %50

Powder for Oral Solution

500 mg/g Neomycin

TARGET SPECIES: Calves, Lambs, Kids, Chicken, Turkey

INDICATIONS: It is indicated for the treatment of infections caused by neomycin sensitive bacteria in calves, lambs, kids chicken and turkey.

PACKAGING: 20 g, 1 kg

DOSE: 1 gr/ 20 kg b.w./ day, split in 2 for lambs, calves and kids for 3-5 days

WITHDRAWAL TIME:

Meat: 14 days

Egg: zero "0" days

See the leaflet for detailed information.



Paripto

Powder for Oral Solution

100 mg/g Paromomycine Sulfate

TARGET SPECIES: Non-Ruminating Calves

INDICATIONS: Paripto Powder for Oral Solution is used for the treatment of gastro-intestinal infections caused by E. Coli, Salmonella and Cryptosporidium in non-ruminating calves

PACKAGING: 1 kg, 2,5 kg

DOSE: Used by adding to drinking water

2,5-5 g/ 10 kg bw/day for Colibacillosis and Salmonellosis, during 3-5 days. 5 g/ 10 kg bw/day for Cryptosporidiosis during 7 days

WITHDRAWAL TIME: 62 days (5 g/ 10 kg b.w. for 7 days treatment)

Meat: 30 days (2,5-5 g/ 10 kg b.w. for 3-5 days treatment)

See the leaflet for detailed information.



Spiramezol

Oral Tablet

750,000 IU/tb Spiramycin and 125 mg/tb Metronidazole

TARGET SPECIES: Dog

INDICATIONS:

It is effective against gram-positive bacteria and anaerobes sensitive to the spiramycin/ metronidazole combination. In the treatment of multiple bacterial infections of the gums and mouth in dogs - gingivitis, stomatitis, glossitis, periodontitis, tonsillitis, dental fistulas and fistulas due to oral wounds, cheilitis and sinusitis - as an adjunct to mechanical or surgical therapy. It is effective against gram positive bacteria and anaerobes sensitive to the spiramycin/ metronidazole combination.

PACKAGINGS:

10 tablet

DOSE:

The practical dosage is 1 tablet per 10 kg body weight. If the product needs to be divided, it should only be divided in half.

WITHDRAWAL TIME:

It is not used in animals with food value.

See the leaflet for detailed information.



Sulfasym

Oral Suspension

400 mg/ml Sulphametoxazole, 80 mg/ml Trimetophrim

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Sulfasym is indicated for the treatment of respiratory infections caused by trimetoprim- and sulfamethoxazole-sensitive *Escherichia Coli* in avians (broiler chickens - turkeys).

PACKAGING: 100 ml, 1 L, 2.5 L

DOSE: There should be 1 ml Sulfasym Oral Suspension per 10.68 kg body weight. Treatment should be continued for 3 days

WITHDRAWAL TIME:Meat: 10 days Should not given to poultry producing eggs for human consumption

See the leaflet for detailed information.



Symtay

Powder for Oral Solution

1000 mg/g Tylosin Tartrate

TARGET SPECIES: Calf, Chicken, Turkey

INDICATIONS: SYMTAY Powder for Oral Solution is indicated for the treatment of infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD), Leptospirosis and infectious sinus inflammation in turkeys.

PACKAGING: 20 g , 100-500-1000

DOSE:

Calves: 22 - 44 mg/kg (7-14 days)

Turkeys-Chickens: 82.5 - 110 mg/kg (3-5 days)

Chickens (Necrotic Enteritis): 22 mg/kg (3 days)

WITHDRAWAL TIME:

Meat : Calves : 12 days , Turkey 2 days, Chicken 1 days

Egg : 0 days

See the leaflet for detailed information.



Symtil %30

Oral Solution

300 mg/ ml Tilmicosine

TARGET SPECIES: Broiler, Turkey, Calves

INDICATIONS: SYMTAY Oral Solution is indicated for the treatment of infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD), Leptospirosis and infectious sinus inflammation in turkeys.

PACKAGING: 1 Lt

DOSE:

Broilers: 5 - 7 ml / 100 kg (3 days)

Turkeys: 3.5 - 9 ml / 100 kg (3 days)

Calves: 1 ml / 24 kg (twice daily for 3 - 5 days)

WITHDRAWAL TIME: Meat: 12 days (chicken), 19 days (turkey), 42 days (calf)

Not to be used in chicken and turkey producing egg intended for human consumption. Not to be used within 14 days of egg laying period.

See the leaflet for detailed information.



Taylosym %24

Oral Solution

240 mg/ml Tylosin Tartarate

TARGET SPECIES: Chicken, Turkey

INDICATIONS: Taylosym 24% Oral Solution is indicated for the treatment of respiratory infections caused by sensitive bacteria in chickens and turkeys. It is mainly used in Mycoplasmosis (CRD) and infectious sinus inflammation in turkeys.

PACKAGING: 1L ve 2.5 L

DOSE:

Turkeys-Chickens (General/CRD): 34 - 45 ml / 100 kg (3-5 days)

Chickens (Necrotic Enteritis): 9 ml / 100 kg (3 days)

WITHDRAWAL TIME: Meat:: 2 days (turkey) 1 days (chickens)

Egg : 0 days

See the leaflet for detailed information.



Tetrasym %80

Powder for Oral Solution

800 mg/g Oxytetracycline

TARGET SPECIES: Calves, Chicken, Turkey, Lamb, Kids

INDICATIONS: TETRASYM 80% Powder for Oral Solution is indicated for the treatment of infections caused by sensitive bacteria in calves, chicken, turkey, lamb and kids.

PACKAGING: 1 kg

DOSE: 2.5 g/ 100 kg total b.w./ day

WITHDRAWAL TIME: Meat: 7 days.

Egg: zero "0" days.

See the leaflet for detailed information.



Kantel

Solution for Injection

56,80 mg/ml Paraziquantel

TARGET SPECIES: Cats, dogs

INDICATIONS: Treatment of cestod infestations in cats and dogs

PACKAGING: 20 ml

DOSE:

Practically, 0.1 ml is injected for 1 kg
body weight. (1 ml/10 kg b.w.)
(s.c.)(i.m.)

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Midokar

Solution for Injection

120 mg/ml Imidocarb Dipropionate

TARGET SPECIES: Cattle, Dog, Horse

INDICATIONS: It is indicated for the treatment and prevention of babesiosis in cattle, dog and horse.

PACKAGING: 20 ml, 50 ml

DOSE:

Cattle: 1ml/100 kg bw for treatment, 2.5 ml/100 kg bw for prevention (i.m.)(s.c.)

Dog: 0.25 ml/10 kg bw for treatment, 0.5 ml/10 kg bw for prevention (i.m.)(s.c.)

Horse: 2 ml/100 kg bw for treatment and prevention (i.m.)(s.c.)

WITHDRAWAL TIME: Meat: 213 days, Milk: 21 days

See the leaflet for detailed information.



Klosym

Closantel sodium dihydrate equivalent of 50 mg/ml Closantel sodium base

TARGET SPECIES:Cattle, Sheep

INDICATIONS: Treatment of closantel sensitive trematods, nematods and arthropod larvae invasions.

PACKAGING: 100 ml

Solution for Injection

DOSE: 1-2 ml/20 kg b.w., s.c. For cattle, max. 20 ml (6 ml in calves) for an injection area. 1-2 ml/20 kg b.w., s.c. for in sheep, max. 5 ml for an injection area.

WITHDRAWAL TIME: Meat: 84 days (cattle), 107 days (sheep).
Not to be used on animals producing milk for human consumption or cattle at last trimester.

See the leaflet for detailed information.



Symec %1

Solution for Injection

10 mg/ml Ivermectin

TARGET SPECIES: Cattle, Sheep

INDICATIONS: It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, subcutaneous nematodes, eye nematode, noca agents, scabies, lice and tick infestations in cattle.

It is used in the treatment and control of gastro-intestinal nematodes, lung nematodes, nasal nematodes, scabies and lice infestations in sheep.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: Practical dosage is 1 ml for 50 kg bodyweight
Only given subcutaneously.

WITHDRAWAL TIME:Meat: 49 days (cattle), 42 days (sheep) Not to be used within 60 days to birth in animals producing milk intended for human consumption

See the leaflet for detailed information.



Symec Fascio

Solution for Injection

10 mg/ml Ivermectin, 100 mg/ml Clorsulon

TARGET SPECIES: Cattle

INDICATIONS: It is used in the prevention and treatment of gastrointestinal nematodes, lung nematodes (adult and 4th stage larvae), subcutaneous nematode, eye nematode, nocturnal agents (larval stage), lice, scabies and liver trematodes in cattle.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 1 ml for 50 kg/ bodyweight Use subcutaneously only

WITHDRAWAL TIME:

Meat: 66 days,
Not authorised to use on cattle that produce milk for human consumption.
Not to be used within 60 days before giving birth.

See the leaflet for detailed information.



Albesym %10

Oral Suspension

100 mg/ml Albendazole

TARGET SPECIES: Cattle, Sheep, Goat

INDICATIONS: It is indicated to treatment and prevention of infestations caused of gastrointestinal nematode, lungworm and trematodes.

PACKAGING: 1 L

DOSE:

Cattle: 7,5 ml/ 100 kg b.w. for gastrointestinal roundworms and lungworms and 10 ml/ 100 kg b.w. for adult flukes

Sheep: 1 ml/ 20 kg b.w. for gastrointestinal roundworms and lungworms and 1,5 ml/ 20 kg b.w.for adult flukes, Dicrocoeliasis dose is 3 ml/20 kg b.w.

Goats: worm and fluke dose is 1,5 ml/ 20 kg b.w. and Dicrocoeliasis dose is 6 ml/40 kg b.w.

WITHDRAWAL TIME:

Meat: 14 days for cattle, 10 days for sheep and goats

Milk: 3 days for cattle, 6 days for sheep and goats (not for human consumption during treatment)

See the leaflet for detailed information.



Albesym 1200

Oral Tablet

1200 mg/ tb Albendazole

TARGET SPECIES: Cattle

INDICATIONS: It is indicated to treatment and prevention of infestations caused by worms, liver flukes.

PACKAGING: 10 tablets

DOSE: 1 tablet/ 160 kg b.w. for worms and yung flukes and 1 tablet/ 120 kg b.w.for adult flukes

WITHDRAWAL TIME: Meat: 14 days

Milk: 3 days (6 milkings) (not for human consumption during treatment)

See the leaflet for detailed information.



Levasym

Oral Suspension

30 mg/ml Levamisole HCl 60 mg/ml Oxytoclozanide

TARGET SPECIES: Cattle, Sheep

INDICATIONS: It is used in the treatment and control of the following gastrointestinal and pulmonary nematodes and adult liver fluke infestations in cattle and sheep:

PACKAGING: 1L

DOSE:

Cattle: 25 ml per 100 kg body weight

Sheep: 2.5 ml per 10 kg body weight

WITHDRAWAL TIME:

Meat: Cattle: 20 days Sheep: 17 days

Milk: Not for use in animals producing milk for human consumption

See the leaflet for detailed information.



Medizanid

Oral Tablet

300 mg/tb Oxfendazole, 600 mg/tb Oxyclosanide

TARGET SPECIES: Cattle

INDICATIONS: Medizanid Oral Tablet is used in prophylaxis and treatment of gastrointestinal and pulmonary infestation by mature and larvae of pinworm, and mature liver flukes, mature *Moniezia* sp. in cattle, camel and sheep.

- a) Liver Flukes
- b) Gastrointestinal pinworms
Stomach
Intestine
Large intestine
- c) Pulmonary pinworms
- d) Tapeworm

PACKAGING: 10 and 20 tablet

DOSE:

Lamb: $\frac{1}{2}$ tablet for 25 kg

Sheep: 1 tablet for 50 kg

Calf-Steer: 2 tablets for 100 kg

Cattle: 10 tablet for 500 kg

WITHDRAWAL TIME: Meat: 28 days (cattle) , 21 days (sheep)

Milk: 5 days (cattle) Not to be used for dairy cows producing milk for human consumption.

See the leaflet for detailed information.



Prazifen KK

Oral Tablet

50 mg/tb Praziquantel, 500 mg/tb Fenbendazole

TARGET SPECIES: Cat ,Dog

INDICATIONS: Used for the treatment of infestations caused by roundworms and tapeworms in dogs and cats

PACKAGING: 3*10 tablets blister1*50 tablets sachet in cardboard box

DOSE: 1 tablet for 10 kg bodyweightTablets may also be administered half according to the weight of the animals

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Wormkill

50 mg/tb Praziquantel, 144 mg/tb Pirantel pamoat 150 mg/tb Febantel

TARGET SPECIES: Dog

INDICATIONS: WORMKILL Oral Tablet is indicated for the control of and fighting against the parasitic invasions caused by the following nematode and cestode species in dogs. Ascarid: *Toxocara canis*, *Toxascaris leonina* (mature and immature) Hookworm: *Uncinaria stenocephala*, *Ancylostoma caninum* (mature) Whipworm: *Trichuris vulpis* (mature) Cestode: *Echinococcus granulosus*, *Echinococcus multilocularis*, *Dipylidium caninum*, *Taenia* sp.

Oral Tablet

PACKAGING: 50 tablets

DOSE: Practical dosage for dogs: 1 tablet for 10 kg bodyweight Tablets may also be administered half according to the weight of the animals.

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Eprinosym

Pour-On Solution

5 mg/ml Eprinomectin

TARGET SPECIES: Cattle

INDICATIONS: It is used for the treatment and control of gastro intestinal pinworms (adult and larval forms), lung pinworms (adult and larval forms), lice, chorioptic and sarcoptic mange and Haematobia irritans in cattle.

PACKAGING: 500 ml

DOSE: 1 ml/10 kg body weight. It should be applied along the back line from the shoulders to the head of the tail. Its effect lasts 2-4 weeks

WITHDRAWAL TIME: Meat: 10 days, Milk: "0" days

See the leaflet for detailed information.



Flumetsym

Pour on Solution

10 mg/ml Flumethrin

TARGET SPECIES: Cattle, Sheep

INDICATIONS: Flumetsym Pour on Solution is used for the control and treatment against biting flies, chewing and biting lice (Haematopinus eury-sternus, Linognathus vituli, Bovicola bovis, B. ovis, Melophagus ovinus), ticks, mange mites (scabies) and other ecto parasites related infections in cattle, sheep and dogs.

PACKAGING: 100 ml, 500 ml

DOSE: The application dose is 1 mg/kg b.w. for tick control in cattle and sheep, and 2 mg/kg body weight for scabies and lice control.

WITHDRAWAL TIME:

Meat: 21 days (sheep) 5 days (cattle)

Milk: 8 days

Not authorised to use on sheep that produce milk for human consumption.

See the leaflet for detailed information.



Amprolsym %25

Oral Solution

250 mg/ml Amprolium

TARGET SPECIES: Chicken, Turkey


INDICATIONS: Treatment and prevention of coccidiosis caused by Eimeria tenella, E. Necatrix, E. Acervulina, E. Maxima, E. Brunetti in chicken

PACKAGING: 1 L

DOSE: Chickens-Turkeys: 20 mg amprolium/kg (5 - 7 days)

WITHDRAWAL TIME:
"0" days for milk and egg

See the leaflet for detailed information.



Symcox

Oral Solution

25 mg/ml Toltrazuril

TARGET SPECIES: Chicken, Turkey

INDICATIONS: It is indicated for the treatment of and protection against coccidiosis in chickens and turkeys. Below are the most important coccidiosis agents, which are included in the spectrum of action of Symcox Oral Solution.; For chickens: Eimeria acervulina, Eimeria brunetti, Eimeria maxima, Eimeria mivati, Eimeria necatrix, Eimeria tenella For turkeys: Eimeria adenoides, Eimeria meleagritidis

PACKAGING: 100 ml, 1000 ml

DOSE: 280 ml product/1000 kg total body weight

WITHDRAWAL TIME:

Meat: 21 days

It should be administered to chickens and turkeys producing egg for human consumption.

See the leaflet for detailed information.



IPM Ksilazin %2

Solution for Injection
Veterinary Sedative, Hypnotic

20 mg/ml Xylazine

TARGET SPECIES: Horse, Cattle, Dog, Cat

INDICATIONS: For sedation, muscle relaxation and analgesia in minor operations, as well as premedication for general anesthesia

PACKAGING: 50 ml

DOSE: It is administered by intravenous, intramuscular and subcutaneous routes in target species.

Please read leaflet for detailed information

WITHDRAWAL TIME:Meat: 1 day

Milk "0" day

See the leaflet for detailed information.



Furosym

Solution for Injection

20 mg/ml Furosemide

TARGET SPECIES: Cattle, Horse, Cat, Dog

INDICATIONS: Furosym Solution for Injection is used in order for elimination of fluids, accumulating and causing the formation of edema due to the disorders in circulatory system, liver, kidneys and any other reasons, from the tissues, body cavities, joints, tendons and surfaces through the diuresis, and in order for acceleration of diuresis in case of intoxications, and also it is used in case of emergencies such as acute lung and brain edema.

PACKAGING: 20 ml, 50 ml

DOSE: Practical dosage: 2,5 - 5 ml/100 kg/bw for cattle and horses (slowly intravenous), 1,25 -2,5 ml/10 kg/bw for dog and cats (slowly intravenous or intramusculary)

WITHDRAWAL TIME: "0 days" meat and milk (i.v.)

See the leaflet for detailed information.



Symatro

Solution for Injection

2 mg/ml Atropine

TARGET SPECIES: Horse,Cat,Dog

INDICATIONS: For stimulation of hearth movements in horse, dog and cats,
at cases of pain, bladder, uretra and kidney spasms colics and constrictions as spasmolitic,
at organic phosphorus poisoning with aracoline, pilocarpin and neostigmin poisonings as antidote
at general anesthesia for reduction of saliva, for prevention of vago-vagal reflexes and vagal tonus negative effects on hearth as preanesthetic medication.

PACKAGING: 20 ml and 50 ml

DOSE:

25 - 500 µg/kg body weight

WITHDRAWAL TIME: Not applicable

See the leaflet for detailed information.



Symbuton

Solution for Injection

100 mg/ml Menbuton

TARGET SPECIES: Cattle, Horse, Sheep, Goat

INDICATIONS: Stimulation of hepato-digestive activity in case of digestive disorders and hepatic insufficiency


PACKAGING: 100 ml

DOSE:

Cattle : 1 ml for 15-20 kg/ bodyweight by intravenously
 Sheep, goat and calves : 1 ml for 10 mg/kg bodyweight by deep intramuscular injection or intravenous slowly
 Horse : 1 ml for 20-40 kg/ bodyweight by intravenous injection

WITHDRAWAL TIME: '0' days for meat and milk

See the leaflet for detailed information.





Adecure AD₃E

500.000 IU/ml Vitamin A, 75.000 IU/ml Vitamin D₃,
50 mg/ml Vitamin E

TARGET SPECIES: Cattle, Horse Sheep, Goat

INDICATIONS: A, D₃ and E avitaminosis and treatment for related diseases and malnutritions.

Solution for Injection

PACKAGING: 100 ml and 250 ml

DOSE:

Calves: 2 ml (IM)

Cattle, Horses: 10 ml (IM)

Kids-Lambs: 0.5 ml (IM)

Sheep-Goats: 1 ml (IM)

WITHDRAWAL TIME:

Meat: 308 days (Cattle), 215 days (Sheep-Goats)

Milk: 5 days (10 milkings)

See the leaflet for detailed information.



Adezek

Solution for Injection
Vitamin

80 mg/ml Vitamin A (200.000 IU), 0,7875 mg/ml Vitamin D3 (30.000 IU), 20 mg/ml Vitamin E

TARGET SPECIES: Cattle, Sheep, Goat, Rabbit

INDICATIONS: Adezek Solution for Injection is indicated for to prevent deficiencies and treatment of diseases due to avitaminosis of these vitamins in cattle, sheep, goat and rabbits.

PACKAGING: 100 ml

DOSE:

Cattle: 5-10 ml

Calves: 2-4 ml

Sheep- Goat: 1-2 ml

Lamb-Rabbit: 0.5 ml

WITHDRAWAL TIME:

Cattle 271 days, Sheep and Goat : 206 days

See the leaflet for detailed information.



Bevitbar

Solution for Injection

100 mg/ml Vitamin B1, 10 mg/ml Vitamin B6

TARGET SPECIES:Horse,Cattle, Sheep, Goat, Cat, Dog

INDICATIONS:Bevitbar Solution for Injection is used against Vitamin B1 (Thiamin HCl) and Vitamin B6 (Pyridoxine HCl) deficiencies or as supportive to treatment in cattle, sheep, goat, cat and dog.

PACKAGING:100 ml

DOSE:

Cattle - Horse 400 kg 10 -20 ml daily

Calf - Heifer 200 kg 5 – 10 ml daily

Sheep - Goats 100 kg 2,5 - 5 ml daily

Cat - Dog 5 kg 1 -2 ml daily

WITHDRAWAL TIME:"0" days for milk and meat

See the leaflet for detailed information.



Selesym

Solution for Injection

60 mg/ml Vitamin E (Tocopherole acetate), 40 mg/ml Vitamin B1 (Thiamin HCl), 1 mg/ml Sodium Selenite

TARGET SPECIES: Cattle, Horse, Sheep, Goat, Calf, Foal, Lamb, Kid

INDICATIONS: It is indicated for white muscle disease, myopathies, growth retardation due to selenium and vitamin E deficiency, and nutritionally-related fertility loss in lambs and calves, cows, sheep, goats, and foals. It is also administered to animals grazing on selenium-deficient land or those whose diet is deficient in vitamin E to protect their offspring. It is used for certain skin diseases and scurvy in calves and for azoturia resulting from myoglobinemia in horses.

PACKAGING: 100 ml

DOSE:

Species	Body weight (kg)	Prophylaxis	Therapeutic
Cattle, horse (After the 5th month of pregnancy)	100	5-10 ml	-
Sheep, goat (After the 3rd month of pregnancy)	50	1-2 ml	-
Calf, Foal	50	0.5-1 ml	1.2 ml
Lamb, goat (1-7 days)	10	0.25 ml	0.25-0.5 ml
Lamb, goat (7-30 days)	10	0.25-0.5 ml	0.5-1.0 ml
Lamb, goat (older than 1 month)	20	0.5-1.0 ml	1.2 ml

WITHDRAWAL TIME: Meat: 28 days.
Not to be used in animals producing milk intended for human consumption.

See the leaflet for detailed information.



Symvit-C

Solution for Injection

200 mg /ml Vitamin C (Ascorbic acid)

TARGET SPECIES: Cattle, Sheep, Horse, Goat, Cat, Dog

INDICATIONS: It is used to treatment and prevention diseases caused by ascorbic acid (vitamin C) deficiency in animals. It is used in clinical and supportive treatment in cases of developmental and adaptation deficiencies, anemia, hypocalcemia, mycotoxicosis and intoxication

PACKAGING: 50 ml, 100 ml

DOSE:

Cattle, 7 - 12 ml /day i.m , s.c , i.v twice a week during 3-6 weeks

Sheep, 2 - 4 ml/day i.m , s.c twice a week during 3 weeks

Goat, 2 - 4 ml/day i.m , s.c twice a week during 3 weeks

Horse, 10 -20 ml /day i.m, i.v

Cat, 0,3 - 0,6 ml/day i.m, i.v

Dog, 0,3 – 2,5 ml/day i.m, i.v

WITHDRAWAL TIME: "0" days for meat and milk

See the leaflet for detailed information.



Symfos

Solution for Injection

100 mg/ml Butaphosphan, 0.05 mg/ml Vitamin B12
(Cyanocobalamine)

TARGET SPECIES: Cattle, Horse, Cat, Dog

INDICATIONS: It is used as a supplement in muscle, metabolism or reproductive system disorders where the use of phosphorus and cyanobalamin is required in cattle, horse, cats and dogs.

PACKAGING: 50 ml, 100 ml, 250 ml

DOSE: 2-5 ml/ 100 kg b.w. (i.v.) in horses and cattle.

0,35-0,6 ml/10 kg b.w. in foals and calves.

0,25-2,5 ml/ 10 kg b.w. (i.v., i.m., s.c.) in dogs.

0,5-5 ml/ 5 kg b.w. (i.v., i.m., s.c.) in cats.

Doses must be split in half with 1-2 weeks delay for chronic conditions in cats and dogs.

WITHDRAWAL TIME: '0' days for meat and milk (cattle, horse)

See the leaflet for detailed information.



Symcal

Solution for Injection

190 mg/ml Calcium Gluconate, 45 mg/ ml Calcium Glucoheptonate, 60 mg/ml Magnesium Chloride, 4 mg/ml Butaphosphane, 10 mg/ml Calcium D-Saccharide

TARGET SPECIES: Cattle, Horse, Calves, Foals, Sheep, Goat, Dog

INDICATIONS: It is indicated for the protection from and treatment of acute and chronic metabolism disorders resulted from the deficiency of calcium, magnesium and phosphor in cattle, horses, sheep, goats and dogs.

PACKAGING: 100 ml, 250 ml

DOSE: Use intramuscularly, intravenously or subcutaneously
Practical dosage: 1 ml/1 kg bw

WITHDRAWAL TIME: "0" days for meat and milk (cattle, sheep, goat)

See the leaflet for detailed information.



Badeljin

Solution for Injection

500 mg/ml Metamisole Sodium

TARGET SPECIES: Cattle, Horse, Dog

INDICATIONS: Badeljin Solution for Injection is used in the symptomatic treatment of the following cases in cattle, horses and dogs.

As analgesic, Spasmolithically, Antipyretically, Antirheumatic

PACKAGING: 50 ml and 100 ml

DOSE:

Horse - Cattle: 4 - 10 ml / 100 kg

Foal - Calves: 2 - 6 ml / 40 kg

Dogs: 0.5 - 1 ml / 10 kg

Horses: Slow intravenous (IV) injection only.

Other Animals: Deep intramuscular (IM) injection or slow IV for severe pain.

WITHDRAWAL TIME: Meat : 18 days (Cattle)

Milk : 2 days (Cattle)

See the leaflet for detailed information.



Ketozek

100 mg/ml Ketoprofen

TARGET SPECIES: Cattle, Horse

INDICATIONS: Diseases affecting the osteoarticular and muscular-skeletal system associated with acute pain and inflammation, after calving.

PACKAGING: 100 ml and 250 ml

Solution for Injection
Non-steroidal Anti-inflammatory, Antiromatizmal

DOSE:

For Cattle 3 ml/ 100 kg b.w. use intramuscular or intravenously, for up to 3 days

For horse 1 ml/ 45 kg b.w. for horse use intravenously, for 3-5 days.

WITHDRAWAL TIME: Meat: 4 days **Milk:** Zero days.

See the leaflet for detailed information.



Melosym

Solution for Injection

5 mg /ml Meloxicam

TARGET SPECIES: Cattle,Dog,Cat

INDICATIONS: In cattle; It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory tract infection. In dogs; It is used to relieve inflammation and relieve pain in acute and chronic musculoskeletal diseases, and to reduce inflammation and pain after orthopedic and soft tissue operations. in cats; It is used to reduce post-operative pain after ovariohysterectomy and mild soft tissue operations.

PACKAGING: 20 ml, 50 ml, 100 ml

DOSE: Cattle: 10 ml/100 kg bw subcutaneously or intravenously

Dogs: 04 ml/10 kg bw subcutaneously

Cat: 0.12ml/2 kg bw subcutaneously

WITHDRAWAL TIME:

Meat: 15 days (Cattle) **Milk:** 5 days (10 milkings) (cattle)

See the leaflet for detailed information.



Melosym X8

40 mg/ml Meloxicam

TARGET SPECIES: Cattle, Horse

INDICATIONS: It is used together with appropriate antibiotics to reduce the clinical manifestations of acute respiratory infection.

PACKAGING: 20 ml and 100 ml

Solution for Injection

DOSE:

Cattle: 1.25 ml / 100 kg b.w. by single dose subcutaneous (sc) or intravenous (iv) injection.

Horse: It is administered by a single dose intravenous (iv) i 1.5 ml / 100 kg b.w.

WITHDRAWAL TIME:

Meat: 15 days (Cattle) **Milk:** 5days (10 milkings) (cattle)

See the leaflet for detailed information.



Symful

Solution for Injection

50 mg/ ml Flunixin

TARGET SPECIES: Cattle, Horse

INDICATIONS: Horse: Painful and inflammatory conditions of the skeletal-muscular system and colic.

Cattle: Accompanying proper treatment, to reduce clinical symptoms formed during pulmonary infections.

DOSE: Intravenous application should be slow and administered at body temperature.

2 ml/ 50 kg b.w. (slow i.v., i.m.) for 1-3 days in cattle.

1 ml/ 50 kg b.w. (slow i.v.) for 1-5 days in horses.

WITHDRAWAL TIME:Meat: 31 days (i.m.), 10 days (i.v.)

Milk: 36 hours (i.m.), 24 hours (i.v.)

See the leaflet for detailed information.

PACKAGING: 100 ml, 250 ml





Tepeören İtosh Mah. 3. Cad. No:7 Tuzla - İstanbul

T: +90 0216 290 22 44 info@ipmilac.com

www.ipmilac.com

